

Derivation of multi-fluid models

MODTERCOM

Matthieu Hillairet (I3M, Université Montpellier 2)

Results published in :

M.H., *JMFM* '07

D. Bresch & X. Huang, *ARMA* '14

D. Bresch & M.H. '14

Compressible multiphase flows

"Definition" [A. Murrone, PhD '04]

Mixture of several compressible phases at equilibrium in which the topology, composition and transfers are parameters that may constantly vary in time and space.

Comestible examples : soda, champagne, emulsions

Applications :

- Simulating aerated flows
(Wave breaking)
- Nuclear industry



Modeling of compressible multiphase flows

Compressible multiphase flows

Modeling [M. Ishii '75, D. Drew & S. Passman '98]

Phase variables :

density ρ_k , velocity u_k , pressure p_k , strain tensor τ_k

Constitutive equations for phase k ($= +, -$)

$$\left\{ \begin{array}{l} \partial_t \rho_k + \operatorname{div}(\rho_k u_k) = 0 \\ \partial_t(\rho_k u_k) + \operatorname{div}(\rho_k u_k \otimes u_k) = \operatorname{div} \tau_k - \nabla p_k \\ \rho_k = \mathcal{P}_k(\rho_k). \end{array} \right. \quad \text{on } \mathcal{F}_k(t)$$

Extended constitutive equations : $X_k = \mathbf{1}_{\mathcal{F}_k}$, velocity σ

$$\left\{ \begin{array}{l} \partial_t(\rho_k X_k) + \operatorname{div}(\rho_k u_k X_k) = \rho_k (u_k - \sigma) \cdot \nabla X_k \\ \partial_t(\rho_k u_k X_k) + \operatorname{div}(X_k \rho_k u_k \otimes u_k) = \operatorname{div}(X_k \tau_k) - \nabla X_k p_k \\ \quad + \rho_k (u_k - \sigma) \otimes u_k \nabla X_k + (p_k \mathbb{I} - \tau_k) \nabla X_k \\ \rho_k = \mathcal{P}_k(\rho_k). \end{array} \right.$$

Compressible multiphase flows

Modeling [M. Ishii '75, D. Drew & S. Passman '98]

Mean operator : $\langle \cdot \rangle$

$$\alpha_k = \langle X_k \rangle \quad \bar{\rho}_k = \frac{\langle X_k \rho_k \rangle}{\langle X_k \rangle} \quad \bar{p}_k = \frac{\langle X_k p_k \rangle}{\langle X_k \rangle} \quad \bar{\tau}_k = \frac{\langle X_k \tau_k \rangle}{\langle X_k \rangle} \quad \tilde{u}_k = \frac{\langle X_k \rho_k u_k \rangle}{\langle X_k \rho_k \rangle}.$$

Homogenized system for phase k ($= +, -$)

$$\left\{ \begin{array}{l} \partial_t(\alpha_k \bar{\rho}_k) + \text{div}(\alpha_k \bar{\rho}_k \tilde{u}_k) = \Gamma_k \\ \partial_t(\alpha_k \bar{\rho}_k \tilde{u}_k) + \text{div}(\alpha_k \bar{\rho}_k \tilde{u}_k \otimes \tilde{u}_k) = \text{div}(\alpha_k (\bar{\tau}_k + \tau_k^T)) - \nabla(\alpha_k \bar{p}_k) \\ \quad + M_k^\Gamma + p_k^{\text{int}} \nabla \alpha_k + F_k \\ \bar{p}_k = \bar{P}_k(\bar{\rho}_k) + p_k^T. \end{array} \right.$$

where :

$$\Gamma_k = \langle \rho_k (u_k - \sigma) \cdot \nabla X_k \rangle, \quad F_k = \langle \tau_k \nabla X_k \rangle, \quad \dots$$

Baer-Nunziato models

Algebraic closure

Modeling assumptions :

$$\Gamma_k = 0, \quad M_k^\Gamma = 0, \quad F_k = \frac{1}{\mu}(u_{k'} - u_k) \quad \tau_k^T = 0 \quad p_k^T = 0.$$

Closure law : $p_+ = p_-$

System :

$$\left\{ \begin{array}{l} \partial_t(\alpha_k \rho_k) + \operatorname{div}(\alpha_k \rho_k u_k) = 0 \\ \partial_t(\alpha_k \rho_k u_k) + \operatorname{div}(\alpha_k \rho_k u_k \otimes u_k) + \alpha_k \nabla p = \frac{1}{\mu}(u_{k'} - u_k) + (p_k^{int} - p) \nabla \alpha_k \\ p = \mathcal{P}_k(\rho_+) = \mathcal{P}_k(\rho_-). \end{array} \right.$$

- $0 \leq \alpha_{\pm}$,
- $\alpha_+ + \alpha_- = 1$

Baer-Nunziato models

Algebraic closure

Modeling assumptions :

$$\Gamma_k = 0, \quad M_k^\Gamma = 0, \quad F_k = \frac{1}{\mu}(u_{k'} - u_k) \quad \tau_k^T = 0 \quad p_k^T = 0.$$

Closure law : $p_+ = p_-$

System :

$$\left\{ \begin{array}{l} \partial_t(\alpha_+ \rho_+) + \operatorname{div}(\alpha_+ \rho_+ \mathbf{u}) = 0 \\ \partial_t \rho + \operatorname{div}(\rho \mathbf{u}) = 0 \\ \partial_t(\rho \mathbf{u}) + \operatorname{div}(\rho \mathbf{u} \otimes \mathbf{u}) + \nabla p = 0 \\ p = \bar{P}_+(\rho_+) = \bar{P}_-(\rho_-). \end{array} \right.$$

with

- $0 \leq \alpha_\pm$,
- $\alpha_+ + \alpha_- = 1$
- $\rho = \alpha_+ \rho_+ + \alpha_- \rho_-$

Baer-Nunziato models

PDE closure

Modeling assumptions :

$$\Gamma_k = 0, \quad M_k^\Gamma = 0, \quad F_k = \frac{1}{\mu}(u_{k'} - u_k) \quad \tau_k^T = 0 \quad p_k^T = 0.$$

$$\text{Closure law : } \partial_t \alpha_+ + u^{int} \cdot \nabla \alpha_+ = \frac{1}{\lambda}(p_+ - p_-)$$

System :

$$\left\{ \begin{array}{l} \partial_t(\alpha_k \rho_k) + \text{div}(\alpha_k \rho_k u_k) = 0 \\ \partial_t(\alpha_k \rho_k u_k) + \text{div}(\alpha_k \rho_k u_k \otimes u_k) + \nabla(\alpha_k p_k) = \frac{1}{\mu}(u_{k'} - u_k) + p_k^{int} \nabla \alpha_k \\ p_k = P_k(\rho_k). \end{array} \right.$$

- $0 \leq \alpha_{\pm},$
- $\alpha_+ + \alpha_- = 1$

Baer-Nunziato models

PDE closure

Modeling assumptions :

$$\Gamma_k = 0, \quad M_k^\Gamma = 0, \quad F_k = \frac{1}{\mu}(u_{k'} - u_k) \quad \tau_k^T = 0 \quad p_k^T = 0.$$

$$\text{Closure law : } \partial_t \alpha_+ + u \cdot \nabla \alpha_+ = \frac{1}{\lambda}(\rho_+ - \rho_-)$$

System :

$$\left\{ \begin{array}{l} \partial_t(\alpha_+ \rho_+) + \operatorname{div}(\alpha_+ \rho_+ u) = 0 \\ \partial_t \rho + \operatorname{div}(\rho u) = 0 \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \nabla p = 0 \\ p = \alpha_+ \mathcal{P}_+(\rho_+) + \alpha_- \mathcal{P}_-(\rho_-). \end{array} \right.$$

with

- $0 \leq \alpha_{\pm},$
- $\alpha_+ + \alpha_- = 1$
- $\rho = \alpha_+ \rho_+ + \alpha_- \rho_-$

Analytical derivation

Composite problem

Composite unknowns :

$$\rho = \rho_+ X_+ + \rho_-(1 - X_+) \quad u = u_+ X_+ + u_-(1 - X_+) \quad p = p_+ X_+ + p_-(1 - X_+).$$

Composite systems :

$$(NS) \quad \begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0 \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \nabla p = \operatorname{div} \tau \end{cases} \quad \text{on } (0, T) \times \Omega$$


Constitutive equations :

$$(CI1) \quad \tau = 2\mu D(u) + \lambda \operatorname{div} u \mathbb{I}$$

$$(CI2) \quad p = \mathcal{P}(\rho)$$

Boundary conditions :

$$u = 0 \quad \text{on } \partial\Omega.$$



Composite pressure diagram

New statement of our problem

Cauchy problem

System (CNS) = (NS) +(CI1)+(CI2) + (BC) is completed with initial condition :

$$(IC) \quad \begin{cases} \rho(0, x) & = & \rho^0(x) \\ u(0, x) & = & u^0(x) \end{cases} \quad \text{on } \Omega$$

where u^0 is given and

$$\rho^0(x) = X_+ \left(x, \frac{x}{\varepsilon} \right) \rho_+^0(x) + \left(1 - X_+ \left(x, \frac{x}{\varepsilon} \right) \right) \rho_-^0(x) \quad \varepsilon \ll 1.$$

Open question :

Given initial data $(\rho_\varepsilon^0, u_\varepsilon^0)_{\varepsilon \rightarrow 0}$, of the above form, and $(\rho_\varepsilon, u_\varepsilon)_{\varepsilon \rightarrow 0}$ the associated solutions to (CNS)+(IC), can we :

- recover (ρ_+, ρ_-) , and (u_+, u_-) ?
- compute equations satisfied by these unknowns ?

1D formal approach

System

$$\begin{aligned}\partial_t \rho + \partial_x(\rho u) &= 0 \\ \rho(\partial_t u + u \partial_x u) &= \mu \partial_{xx} u - \partial_x p \\ p &= \mathcal{P}(\rho)\end{aligned}$$

Ansatz

$$\begin{aligned}\rho(t, x) &= \alpha_+ \left(t, \frac{t}{\varepsilon}, x, \frac{x}{\varepsilon} \right) \rho_+(t, x) + \alpha_- \left(t, \frac{t}{\varepsilon}, x, \frac{x}{\varepsilon} \right) \rho_-(t, x) \\ u(t, x) &= u_0(t, x) + \varepsilon u_1 \left(t, \frac{t}{\varepsilon}, x, \frac{x}{\varepsilon} \right) + \dots\end{aligned}$$

with

$$\begin{aligned}\alpha_{\pm}(t, s, x, y) &\in \{0, 1\}, \quad \bar{u}_1 := \int_{Cell} u_1 dy = 0. \\ Im(\rho_+) \cap Im(\rho_-) &= \emptyset\end{aligned}$$

Identifying powers of ε

Momentum equation :

$$\varepsilon^{-1} \quad \partial_y(\mu \partial_y u_1 - p) = 0$$

$$\varepsilon^0 \quad \bar{\rho}(\partial_t u_0 + u_0 \partial_x u_0) = \mu \partial_{xx} u_0 - \partial_x \bar{p}$$

with :

$$\bar{\rho} = \bar{\alpha}_+ \rho_+ + \bar{\alpha}_- \rho_-$$

$$\bar{p} = \overline{\mathcal{P}(\alpha_+ \rho_+ + \alpha_- \rho_-)} = \bar{\alpha}_+ \mathcal{P}(\rho_+) + \bar{\alpha}_- \mathcal{P}(\rho_-).$$

Closure equation = "Continuity equation * $\beta'(\rho)$ " :

$$\partial_t \bar{\alpha}_+ + u_0 \partial_x \bar{\alpha}_+ = \overline{\alpha_+ \partial_y u_1}.$$

Identifying powers of ε

Momentum equation :

$$\varepsilon^{-1} \quad \mu \partial_y u_1 = p - \bar{p}$$

$$\varepsilon^0 \quad \bar{\rho}(\partial_t u_0 + u_0 \partial_x u_0) = \mu \partial_{xx} u_0 - \partial_x \bar{p}$$

with :

$$\bar{\rho} = \bar{\alpha}_+ \rho_+ + \bar{\alpha}_- \rho_-$$

$$\bar{p} = \overline{\mathcal{P}(\alpha_+ \rho_+ + \alpha_- \rho_-)} = \bar{\alpha}_+ \mathcal{P}(\rho_+) + \bar{\alpha}_- \mathcal{P}(\rho_-).$$

Closure equation = "Continuity equation * $\beta'(\rho)$ " :

$$\partial_t \bar{\alpha}_+ + u_0 \partial_x \bar{\alpha}_+ = \overline{\alpha_+ \partial_y u_1}.$$

Identifying powers of ε

Momentum equation :

$$\varepsilon^{-1} \quad \mu \partial_y u_1 = p - \bar{p}$$

$$\varepsilon^0 \quad \bar{\rho}(\partial_t u_0 + u_0 \partial_x u_0) = \mu \partial_{xx} u_0 - \partial_x \bar{p}$$

with :

$$\bar{\rho} = \overline{\alpha_+ \rho_+} + \overline{\alpha_- \rho_-}$$

$$\bar{p} = \overline{\mathcal{P}(\alpha_+ \rho_+ + \alpha_- \rho_-)} = \overline{\alpha_+} \mathcal{P}(\rho_+) + \overline{\alpha_-} \mathcal{P}(\rho_-).$$

Closure equation = "Continuity equation * $\beta'(\rho)$ " :

$$\partial_t \overline{\alpha_+} + u_0 \partial_x \overline{\alpha_+} = \overline{\alpha_+ \partial_y u_1}.$$

with

$$\overline{\alpha_+ \partial_y u_1} = \frac{\overline{\alpha_+} \overline{\alpha_-}}{\mu} (\mathcal{P}(\rho_+) - \mathcal{P}(\rho_-))$$

Identifying powers of ε

Momentum equation :

$$\begin{aligned}\varepsilon^{-1} \quad & \mu \partial_y u_1 = p - \bar{p} \\ \varepsilon^0 \quad & \bar{\rho}(\partial_t u_0 + u_0 \partial_x u_0) = \mu \partial_{xx} u_0 - \partial_x \bar{p}\end{aligned}$$

with :

$$\begin{aligned}\bar{\rho} &= \overline{\alpha_+ \rho_+ + \alpha_- \rho_-} \\ \bar{p} &= \overline{\mathcal{P}(\alpha_+ \rho_+ + \alpha_- \rho_-)} = \overline{\alpha_+} \mathcal{P}(\rho_+) + \overline{\alpha_-} \mathcal{P}(\rho_-).\end{aligned}$$

Closure equation = "Continuity equation * $\beta'(\rho)$ " :

$$\partial_t \overline{\alpha_+} + u_0 \partial_x \overline{\alpha_+} = \frac{\overline{\alpha_+} \overline{\alpha_-}}{\mu} (\mathcal{P}(\rho_+) - \mathcal{P}(\rho_-)).$$

with

$$\overline{\alpha_+ \partial_y u_1} = \frac{\overline{\alpha_+} \overline{\alpha_-}}{\mu} (\mathcal{P}(\rho_+) - \mathcal{P}(\rho_-))$$

Multi-D approach

On compactness of solutions to (CNS)

References : D. Serre '91, P.-L. Lions, '98,
E. Feireisl & H. Petzeltova, '00, E. Feireisl '01'02

Question : Let $(\rho_\varepsilon, u_\varepsilon)_{\varepsilon \rightarrow 0}$ be a sequence of solution to (CNS) on $(0, T)$ such that for arbitrary ε :

$$(\text{Diss}_\varepsilon) \quad \sup_{t \in (0, T)} \left\{ \int_\Omega \left[\frac{1}{2} \rho_\varepsilon |u_\varepsilon|^2 + Q(\rho_\varepsilon) \right] \right\} + \int_0^T \int_\Omega \mu |\nabla u_\varepsilon|^2 + \lambda |\text{div} u_\varepsilon|^2 \leq M$$

Can we extract a limit (ρ, u) solution to (CNS) ?

Remarks

- $(\text{Diss}_\varepsilon)$ means that we have a sequence of bounded-energy solutions.
- Q is defined by $(Q(z)/z)' = \mathcal{P}(z)/z^2$. In particular,

$$\mathcal{P}(z) = az^\gamma \implies Q(z) = \frac{a}{\gamma - 1} z^\gamma.$$

First issue

Is the limit pressure a function ?

Supplementary assumptions :

- Ω is smooth and bounded
- $\mu > 0$ and $\lambda + 2\mu/3 \geq 0$

Uniform integrability I : from the energy estimate

- u_ε uniformly bounded in $L^\infty(0, T; L^2(\Omega)) \cap L^2((0, T); H_0^1(\Omega))$
- ρ_ε uniformly bounded in $L^\infty(0, T; L^\gamma(\Omega))$

Multiply momentum equation with \mathcal{B}_θ solution to

$$\begin{cases} \operatorname{div} \mathcal{B}_\theta &= \rho^\theta - \frac{1}{|\Omega|} \int_\Omega \rho^\theta & \text{in } \Omega \\ \mathcal{B}_\theta \cdot n &= 0, & \text{on } \partial\Omega. \end{cases}$$

Uniform integrability II :

- ρ_ε uniformly bounded in $L^{\alpha_\gamma}((0, T) \times \Omega)$ with $\alpha_\gamma = \gamma + \frac{2}{3}\gamma - 1$.

Second issue

Can we write an equation for the limit pressure ?

Existence/properties of a weak limit

- $u_\varepsilon \rightharpoonup u$ in $L^2((0, T); H_0^1(\Omega)) - w$ and $L^\infty((0, T); L^2(\Omega)) - w^*$
- $\rho_\varepsilon \rightharpoonup \rho$ in $L^{\alpha\gamma}((0, T) \times \Omega) - w$ and $L^\infty((0, T); L^\gamma) - w^*$
- $\rho_\varepsilon \rightharpoonup \rho$ in $L^{\alpha\gamma/\gamma}((0, T) \times \Omega) - w$

solution to

$$\begin{aligned}\partial_t \rho + \operatorname{div}(\rho u) &= 0 \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \nabla p &= \operatorname{div}(2\mu D(u) + \lambda \operatorname{div} u \mathbb{I})\end{aligned}$$

Difficulty : Recover $p = \mathcal{P}(\rho)$

Alternative method : Obtaining an equation for p

$$\partial_t \rho_\varepsilon + \operatorname{div} \rho_\varepsilon u_\varepsilon = 0$$

Second issue

Can we write an equation for the limit pressure ?

Existence/properties of a weak limit

- $u_\varepsilon \rightharpoonup u$ in $L^2((0, T); H_0^1(\Omega)) - w$ and $L^\infty((0, T); L^2(\Omega)) - w^*$
- $\rho_\varepsilon \rightharpoonup \rho$ in $L^{\alpha\gamma}((0, T) \times \Omega) - w$ and $L^\infty((0, T); L^\gamma) - w^*$
- $\rho_\varepsilon \rightharpoonup \rho$ in $L^{\alpha\gamma/\gamma}((0, T) \times \Omega) - w$

solution to

$$\begin{aligned}\partial_t \rho + \operatorname{div}(\rho u) &= 0 \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \nabla p &= \operatorname{div}(2\mu D(u) + \lambda \operatorname{div} u \mathbb{I})\end{aligned}$$

Difficulty : Recover $p = \mathcal{P}(\rho)$

Alternative method : Obtaining an equation for p

$$\partial_t \beta(\rho_\varepsilon) + \operatorname{div}(\beta(\rho_\varepsilon) u_\varepsilon) + (\beta'(\rho_\varepsilon) \rho_\varepsilon - \beta(\rho_\varepsilon)) \operatorname{div} u_\varepsilon = 0$$

for all $\beta : [0, \infty) \rightarrow \mathbb{R}$

Second issue

Can we write an equation for the limit pressure ?

Existence/properties of a weak limit

- $u_\varepsilon \rightharpoonup u$ in $L^2((0, T); H_0^1(\Omega)) - w$ and $L^\infty((0, T); L^2(\Omega)) - w^*$
- $\rho_\varepsilon \rightharpoonup \rho$ in $L^{\alpha\gamma}((0, T) \times \Omega) - w$ and $L^\infty((0, T); L^\gamma) - w^*$
- $p_\varepsilon \rightharpoonup p$ in $L^{\alpha\gamma/\gamma}((0, T) \times \Omega) - w$

solution to

$$\begin{aligned}\partial_t \rho + \operatorname{div}(\rho u) &= 0 \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \nabla p &= \operatorname{div}(2\mu D(u) + \lambda \operatorname{div} u \mathbb{I})\end{aligned}$$

Difficulty : Recover $p = \mathcal{P}(\rho)$

Alternative method : Obtaining an equation for p

$$\partial_t \bar{\beta} + \operatorname{div}(\bar{\beta} u_\varepsilon) + \overline{(\beta'(\rho_\varepsilon) \rho_\varepsilon - \beta(\rho_\varepsilon)) \operatorname{div} u_\varepsilon} = 0$$

for all $\beta : [0, \infty) \rightarrow \mathbb{R}$ when $\varepsilon \rightarrow 0$.

Convention : $\bar{\cdot} = \lim_{\varepsilon \rightarrow 0} \cdot_\varepsilon$

Second issue

Can we write an equation for the limit pressure ?

Further compactness properties

- The divergence of the momentum equation reads :

$$\Delta((\lambda + 2\mu)\operatorname{div}u_\varepsilon - \mathcal{P}(\rho_\varepsilon)) = \operatorname{div}(\rho_\varepsilon(\partial_t u_\varepsilon + u_\varepsilon \cdot \nabla u_\varepsilon))$$

- **Lemma [E. Feireisl, P.L. Lions]** Given $\beta : [0, \infty) \mapsto \mathbb{R}^+$ then, for arbitrary $\varphi \in C_c^\infty((0, T) \times \Omega)$ there holds :

$$\lim_{\varepsilon \rightarrow 0} \int_0^T \int_\Omega ((\lambda + 2\mu)\operatorname{div}u_\varepsilon - \mathcal{P}(\rho_\varepsilon))\beta(\rho_\varepsilon)\varphi = \int_0^T \int_\Omega ((\lambda + 2\mu)\operatorname{div}u - \mathcal{P}(\rho))\beta(\rho)\varphi$$

Conclusion : There holds :

$$\partial_t \bar{\beta} + \operatorname{div}(\bar{\beta}u) + (\overline{\beta'\rho - \beta})\operatorname{div}u = \frac{(\overline{\beta'\rho - \beta})\rho - \overline{(\beta'\rho - \beta)\rho}}{\lambda + 2\mu}$$

for all $\beta : [0, \infty) \rightarrow \mathbb{R}$

Conclusion

Construction of composite unknowns :

We define $\nu^\varepsilon(t, x, \xi) \in \mathcal{M}_+((0, T) \times \Omega \times [0, \infty))$ s.t. :

$$\langle \nu^\varepsilon, \beta(\xi) \otimes \varphi(t, x) \rangle = \int_0^T \int_\Omega \beta(\rho_\varepsilon(t, x)) \varphi(t, x) dt dx, \text{ a.e..}$$

Then $\nu^\varepsilon \rightarrow \nu$ s.t.

$$\int_0^\infty \beta(z) d\nu(z) = \bar{\beta}.$$

Full system :

$$\partial_t \rho + \operatorname{div}(\rho u) = 0$$

$$\partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \nabla p = \operatorname{div}(2\mu D(u) + \lambda \operatorname{div} u \mathbb{I})$$

$$\partial_t \nu + u \cdot \nabla_x \nu = \frac{1}{\lambda + 2\mu} \partial_\xi [\xi \nu \{G - \mathcal{P}(\xi)\}] + \frac{1}{\lambda + 2\mu} \nu (\rho - \mathcal{P}(\xi)),$$

with :

$$G = (\lambda + 2\mu) \operatorname{div}_x u - p \quad p = \int_0^\infty \mathcal{P}(z) d\nu(z) \quad \rho = \int_0^\infty z d\nu(z).$$

Back to the homogenization problem

Composite system (HCNS)

Remark : With initial data $\rho^0 = X_+(x, x/\varepsilon)\rho_+(x) + (1 - X_+(x, x/\varepsilon))\rho_-(x)$ we have :

$$\nu_{0,x} = \alpha_+(x)\delta_{\rho_+(x)} + (1 - \alpha_+(x))\delta_{\rho_-(x)}, \quad \alpha_+(x) = \frac{1}{|\text{cell}|} \int_{\text{cell}} X_+(x, y) dy$$

Assumption : $\nu = \alpha_+\delta_{\rho_+} + \alpha_-\delta_{\rho_-}$ with $\rho_+(t, x) \neq \rho_-(t, x)$.

$$\partial_t \alpha_+ + u \cdot \nabla \alpha_+ = \frac{\alpha_+(\mathcal{P}(\rho_+) - \rho)}{\lambda + 2\mu}$$

$$\alpha_+(\partial_t \rho_+ + \text{div}(\rho_+ u)) = \alpha_+ \frac{\rho_+(\rho - \mathcal{P}(\rho_+))}{\lambda + 2\mu}$$

$$\partial_t \rho + \text{div} \rho u = 0$$

$$\partial_t(\rho u) + \text{div}(\rho u \otimes u) + \nabla p = \text{div}(2\mu D(u) + \lambda \text{div} u \mathbb{I})$$

where :

- $0 \leq \alpha_{\pm}$ and $\alpha_+ + \alpha_- = 1$
- $\rho = \alpha_+\rho_+ + \alpha_-\rho_-$, and $p = \alpha_+\mathcal{P}(\rho_+) + \alpha_-\mathcal{P}(\rho_-)$.

Back to the homogenization problem

Composite system (HCNS)

Remark : With initial data $\rho^0 = X_+(x, x/\varepsilon)\rho_+(x) + (1 - X_+(x, x/\varepsilon))\rho_-(x)$ we have :

$$\nu_{0,x} = \alpha_+(x)\delta_{\rho_+(x)} + (1 - \alpha_+(x))\delta_{\rho_-(x)}, \quad \alpha_+(x) = \frac{1}{|\text{cell}|} \int_{\text{cell}} X_+(x, y) dy$$

Assumption : $\nu = \alpha_+\delta_{\rho_+} + \alpha_-\delta_{\rho_-}$ with $\rho_+(t, x) \neq \rho_-(t, x)$.

$$\partial_t(\alpha_+\rho_+) + \text{div}(\alpha_+\rho_+u) = 0$$

$$\partial_t\rho + \text{div}\rho u = 0$$

$$\partial_t(\rho u) + \text{div}(\rho u \otimes u) + \nabla p = 0$$

$$p = \mathcal{P}_+(\rho_+) = \mathcal{P}_-(\rho_-)$$

where :

- $0 \leq \alpha_{\pm}$ and $\alpha_+ + \alpha_- = 1$
- $\rho = \alpha_+\rho_+ + \alpha_-\rho_-$,

Results

Cauchy theory for (CNS) + (IC)

Weak solutions : P.-L. Lions '98, E. Feireisl, A. Novotný & H. Petzeltová '01

Theorem. Assume Ω is smooth and bounded and

- $\mathcal{P}(z) = az^\gamma$ with $\gamma > 3/2$ and $a > 0$
- $\mu > 0$ and $\lambda + 2\mu/3 > 0$

Then, given a positive $\rho^0 \in L^\gamma(\Omega)$ and $q^0 (= \rho^0 u^0)$ compatible with ρ^0 there exists a finite-energy weak solution (ρ, u) to (CNS)+(IC) on arbitrary large times.

Semi-strong solutions : D. Hoff '95, B. Desjardins '97,

Theorem. Assume $\Omega = \mathbb{T}^3$. Given a positive $\rho^0 \in L^\infty(\Omega)$ and $u^0 \in H^1(\mathbb{T}^3)$ there exists $T_0 > 0$ and a finite-energy weak solution (ρ, u) to (CNS)+(IC) such that :

- $\rho \in L^\infty((0, T_0) \times \mathbb{T}^3) \quad \nabla u \in L^\infty((0, T); L^2(\mathbb{T}^3))$
- $\sqrt{\rho} \partial_t u \in L^2((0, T_0) \times \mathbb{T}^3) \quad Pu \in L^2((0, T_0); H^2(\mathbb{T}^3))$
- $G := (\lambda + 2\mu) \operatorname{div} u - p \in L^2(0, T_0; H^1(\mathbb{T}^3))$

Remark : T_0 depends on $\|\rho_0; L^\infty(\mathbb{T}^3)\|$ and $\|u : H^1(\mathbb{T}^3)\|$ only

Main result

Theorem [D. Bresch & M.H. and D. Bresch & X. Huang '13]

Let initial data $(\rho_n^0, u_n^0) \in L^\infty(\mathbb{T}^3) \times H^1(\mathbb{T}^3)$ satisfy

- $\|\rho_n^0; L^\infty(\mathbb{T}^3)\| + \|u_n^0; H^1(\mathbb{T}^3)\| \leq C$,
- $0 < 1/C \leq \rho_n^0(x)$
- the Young measures ν_n^0 associated with ρ_n^0 converge weakly to

$$\nu^0 = \alpha_+^0(x)\delta_{\rho_+^0(x)} + (1 - \alpha_+^0(x))\delta_{\rho_-^0(x)} \quad \text{on } \Omega.$$

Then, given $\mathcal{P}(z) = az^\gamma$ with $\gamma > 1$

- there exists $T > 0$ and a semi-strong solution (ρ_n, u_n) to (CNS)+(IC) on $(0, T)$
- Up to the extraction of a subsequence

$$\nu_n \rightharpoonup \nu = \alpha_+ \delta_{\rho_+} + (1 - \alpha_+) \delta_{\rho_-}, \quad u_n \rightharpoonup u \quad \rho_n \rightharpoonup \rho$$

- $(\alpha_+, \rho_+, u, \rho)$ is a solution to (HCNS).

Main steps of the proof

Step 1 : Show that in the limit process

- $\operatorname{div} u \in L^1(0, T; L^\infty(\mathbb{T}^3))$
- control the support of ν

Step 2 : Given (u, p) , construct bounded solutions to :

$$\begin{aligned}\partial_t \alpha_k + u \cdot \nabla \alpha_k &= \frac{\alpha_k (\mathcal{P}(\rho_k) - p)}{\lambda + 2\mu} \\ \partial_t \rho_k + \operatorname{div}(\rho_k u) &= \frac{\rho_k (p - \mathcal{P}(\rho_k))}{\lambda + 2\mu}\end{aligned}$$

Step 3 : Given u and p prove weak-strong uniqueness for the Young measure system where G and u are given as above.

Open questions

Consider more complex models

- General pressure laws
- General compressible Navier Stokes (with internal energy)

Include velocity oscillations

⇒ Towards fluid/solid mixture